

Surname		Other Names	
Centre Number		Candidate Number	
Candidate Signature			

Leave blank
-------------

General Certificate of Education  
 June 2003  
 Advanced Subsidiary Examination



**PHYSICS (SPECIFICATION A)**  
**Unit 2 Mechanics and Molecular Kinetic Theory**

**PA02**

Friday 6 June 2003 Afternoon Session

<p><b>In addition to this paper you will require:</b></p> <ul style="list-style-type: none"> <li>• a calculator;</li> <li>• a pencil and a ruler.</li> </ul>
--

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Time allowed: 1 hour

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

**Information**

- The maximum mark for this paper is 50.
- Mark allocations are shown in brackets.
- The paper carries 30% of the total marks for Physics Advanced Subsidiary and carries 15% of the total marks for Physics Advanced.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- In questions requiring description and explanation you will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary where appropriate. The degree of legibility of your handwriting and the level of accuracy of your spelling, punctuation and grammar will also be taken into account.

**Data Sheet**

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Fundamental constants and values				Mechanics and Applied Physics		Fields, Waves, Quantum Phenomena	
Quantity	Symbol	Value	Units				
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$	$v = u + at$		$g = \frac{F}{m}$	
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$	$s = \left(\frac{u+v}{2}\right)t$		$g = -\frac{GM}{r^2}$	
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$	$s = ut + \frac{at^2}{2}$		$g = -\frac{\Delta V}{\Delta x}$	
charge of electron	$e$	$1.60 \times 10^{-19}$	C	$v^2 = u^2 + 2as$		$V = -\frac{GM}{r}$	
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s	$F = \frac{\Delta(mv)}{\Delta t}$		$a = -\frac{(2\pi f)^2 x}{r}$	
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$	$P = Fv$		$v = \pm 2\pi f \sqrt{A^2 - x^2}$	
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$	$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$		$x = A \cos 2\pi ft$	
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$	$\omega = \frac{v}{r} = 2\pi f$		$T = 2\pi \sqrt{\frac{m}{k}}$	
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$	$a = \frac{v^2}{r} = r\omega^2$		$T = 2\pi \sqrt{\frac{l}{g}}$	
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$	$I = \sum mr^2$		$\lambda = \frac{\omega s}{D}$	
the Wien constant	$a$	$2.90 \times 10^{-3}$	m K	$E_k = \frac{1}{2} I\omega^2$		$d \sin \theta = n\lambda$	
electron rest mass	$m_e$	$9.11 \times 10^{-31}$	kg	$\omega_2 = \omega_1 + \alpha t$		$\theta \approx \frac{\lambda}{D}$	
(equivalent to $5.5 \times 10^{-4}u$ )				$\theta = \omega_1 t + \frac{1}{2} \alpha t^2$		$n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$	
electron charge/mass ratio	$e/m_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$	$\omega_2^2 = \omega_1^2 + 2\alpha\theta$		$n_2 = \frac{n_2}{n_1}$	
proton rest mass	$m_p$	$1.67 \times 10^{-27}$	kg	$\theta = \frac{1}{2} (\omega_1 + \omega_2)t$		$\sin \theta_c = \frac{1}{n}$	
(equivalent to 1.00728u)				$T = I\alpha$		$E = hf$	
proton charge/mass ratio	$e/m_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$	<i>angular momentum</i> = $I\omega$		$hf = \phi + E_k$	
neutron rest mass	$m_n$	$1.67 \times 10^{-27}$	kg	$W = T\theta$		$hf = E_1 - E_2$	
(equivalent to 1.00867u)				$P = T\omega$		$\lambda = \frac{h}{p} = \frac{h}{mv}$	
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$	<i>angular impulse</i> = change of <i>angular momentum</i> = $Tt$		$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$	
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$	$\Delta Q = \Delta U + \Delta W$		<b>Electricity</b>	
atomic mass unit	$u$	$1.661 \times 10^{-27}$	kg	$\Delta W = p\Delta V$		$\epsilon = \frac{E}{Q}$	
(1u is equivalent to 931.3 MeV)				$pV^\gamma = \text{constant}$		$\epsilon = I(R + r)$	
<b>Fundamental particles</b>				<i>work done per cycle</i> = area of loop		$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$	
<i>Class</i>	<i>Name</i>	<i>Symbol</i>	<i>Rest energy</i>	<i>input power</i> = calorific value $\times$ fuel flow rate		$R_T = R_1 + R_2 + R_3 + \dots$	
			/MeV	<i>indicated power</i> as (area of $p - V$ loop) $\times$ (no. of cycles/s) $\times$ (no. of cylinders)		$P = I^2 R$	
photon	photon	$\gamma$	0	<i>friction power</i> = indicated power - brake power		$E = \frac{F}{Q} = \frac{V}{d}$	
lepton	neutrino	$\nu_e$	0	<i>efficiency</i> = $\frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$		$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$	
		$\nu_\mu$	0	<i>maximum possible efficiency</i> = $\frac{T_H - T_C}{T_H}$		$E = \frac{1}{2} QV$	
	electron	$e^\pm$	0.510999			$F = BI$	
	muon	$\mu^\pm$	105.659			$F = BQv$	
mesons	pion	$\pi^\pm$	139.576			$Q = Q_0 e^{-t/RC}$	
		$\pi^0$	134.972			$\Phi = BA$	
	kaon	$K^\pm$	493.821				
		$K^0$	497.762				
baryons	proton	$p$	938.257				
	neutron	$n$	939.551				
<b>Properties of quarks</b>							
<i>Type</i>	<i>Charge</i>	<i>Baryon number</i>	<i>Strangeness</i>				
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0				
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0				
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1				
<b>Geometrical equations</b>							
<i>arc length</i> = $r\theta$							
<i>circumference of circle</i> = $2\pi r$							
<i>area of circle</i> = $\pi r^2$							
<i>area of cylinder</i> = $2\pi rh$							
<i>volume of cylinder</i> = $\pi r^2 h$							
<i>area of sphere</i> = $4\pi r^2$							
<i>volume of sphere</i> = $\frac{4}{3}\pi r^3$							

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

### Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

### Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

### Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

### Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

### Electronics

Resistors

Preferred values for resistors (E24)  
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2  
2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2  
6.8 7.5 8.2 9.1 ohms  
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

### Alternating Currents

$$f = \frac{1}{T}$$

### Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left( \frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

Answer **all** questions.

1 A tray containing 0.20 kg of water at 20 °C is placed in a freezer.

(a) The temperature of the water drops to 0 °C in 10 minutes.

specific heat capacity of water = 4200 J kg<sup>-1</sup> K<sup>-1</sup>

Calculate

(i) the energy lost by the water as it cools to 0 °C,

.....  
 .....

(ii) the average rate at which the water is losing energy, in J s<sup>-1</sup>.

.....  
 .....

(3 marks)

(b) (i) Estimate the time taken for the water at 0 °C to turn completely into ice.

specific latent heat of fusion of water =  $3.3 \times 10^5$  J kg<sup>-1</sup>

.....  
 .....

(ii) State any assumptions you make.


.....  
 .....

(3 marks)

6

Turn over ▶

- 2 The diagram represents an experiment that can be used to investigate stopping distances for a moving trolley.



The trolley is placed on the raised section of the track. When released it moves down the track and then travels along the horizontal section before colliding with the block. The trolley and block join and move together after the collision. The distance they move is measured.

- (a) State the main energy changes taking place

(i) as the trolley descends,

.....

(ii) after the collision, as the trolley and block move together.

.....

*(2 marks)*

- (b) Describe how the speed of the trolley, just before it collides with the block may be measured experimentally.

You may be awarded marks for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

*(3 marks)*

(c) State and explain how the speed of the trolley, prior to impact could be varied.

.....  
.....  
.....

(2 marks)

7

3 An apple and a leaf fall from a tree at the same instant. Both apple and leaf start at the same height above the ground but the apple hits the ground first.

You may be awarded marks for the quality of written communication in your answer.

Use Newton's laws of motion to explain why

(i) the leaf accelerates at first then reaches a terminal velocity,

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(ii) the apple hits the ground first.


.....  
.....  
.....  
.....

(5 marks)

5

Turn over ▶

- 4 The diagram shows a 250 kg iron ball being used on a demolition site. The ball is suspended from a cable at point A, and is pulled into the position shown by a rope that is kept horizontal. The tension in the rope is 1200 N.



- (a) In the position shown the ball is in equilibrium.

- (i) What balances the force of the rope on the ball?

.....

- (ii) What balances the weight of the ball?

.....

(2 marks)

- (b) Determine

- (i) the magnitude of the vertical component of the tension in the cable,

.....

.....

- (ii) the magnitude of the horizontal component of the tension in the cable,

.....

.....



(iii) the magnitude of the tension in the cable,

.....  
.....  
.....  
.....  
.....  
.....

(iv) the angle the cable makes to the vertical.

.....  
.....


*(6 marks)*

—  
8

**TURN OVER FOR THE NEXT QUESTION**

**Turn over ▶**

5 The diagram shows the velocity-time graph for a vertically bouncing ball, which is released above the ground at A and strikes the floor at B. The effects of air resistance have been neglected.



(a) (i) What does the gradient of a velocity-time graph represent?

.....

(ii) Explain why the gradient of the line CD is the same as line AB.

.....  
.....

(iii) What does the area between the line AB and the time axis represent?

.....

(iv) State why the velocity at C is negative.

.....  
.....

(v) State why the speed at C is less than the speed at B.

.....  
.....  
.....

(5 marks)

- (b) The ball has a mass of 0.15 kg and is dropped from an initial height of 1.2 m. After impact the ball rebounds to a height of 0.75 m.

Calculate

- (i) the speed of the ball immediately before impact,

.....  
.....

- (ii) the speed of the ball immediately after impact,

.....  
.....

- (iii) the change in momentum of the ball as a result of the impact,

.....  
.....

- (iv) the magnitude of the resultant average force acting on the ball during impact if it is in contact with the floor for 0.10 s.

.....  
.....

(8 marks)

13

**TURN OVER FOR THE NEXT QUESTION**

**Turn over ▶**

- 6 (a) (i) Write down the equation of state for  $n$  moles of an ideal gas.

.....

- (ii) The molecular kinetic theory leads to the derivation of the equation

$$pV = \frac{1}{3}Nmc^2,$$

where the symbols have their usual meaning.

State **three** assumptions that are made in this derivation.

.....

.....

.....

.....

.....

(4 marks)

- (b) Calculate the average kinetic energy of a gas molecule of an ideal gas at a temperature of  $20^\circ\text{C}$ .

.....

.....

.....

.....

(3 marks)

- (c) Two different gases at the same temperature have molecules with different mean square speeds. Explain why this is possible.

.....

.....

.....

.....

(2 marks)

**QUALITY OF WRITTEN COMMUNICATION** (2 marks)

**END OF QUESTIONS**

○  
—  
9

○  
—  
2