

# Mark Scheme Electric Fields Past Paper Questions Jan 2002—Jan 2010 (old spec)

4(a)(i) (force) to the right ✓

(ii) electrons accelerate or speed increases ✓ (2)

Q4 Jan 2002

(b)(i) sketch to show path curving upwards in the field  
(must not become vertical) ✓

(ii) horizontal component of velocity is unchanged ✓  
vertical or upwards acceleration (or force) ✓  
parabolic path described (or named) ✓

max (3)

(5)

3(a)(i)  $E (= \frac{Q}{4\pi\epsilon_0 r^2}) = \frac{29 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{-12} \times (1.15 \times 10^{-10})^2}$  ✓  
 $= 3.15 \times 10^{12} \text{ V m}^{-1} \text{ (or N C}^{-1}\text{)}$  ✓

Q3 Jan 2003

(a)(ii)  $V (= -\frac{GM}{r}) = (-) \frac{6.67 \times 10^{-11} \times 63 \times 1.66 \times 10^{-27}}{1.15 \times 10^{-10}}$  ✓  
 $= (-) 6.07 \times 10^{-26}$  ✓ – sign and  $\text{J kg}^{-1}$  ✓

(5)

(b) arrow pointing to the right ✓

(1)

(6)

2

(a)

Q2 Jan 2004

quantity	SI unit	
(gravitational potential)	$\text{J kg}^{-1}$ or $\text{N m kg}^{-1}$	scalar
(electric field strength)	$\text{N C}^{-1}$ or $\text{V m}^{-1}$	vector
(magnetic flux density)	$\text{T}$ or $\text{Wb m}^{-2}$ or $\text{N A}^{-1} \text{m}^{-1}$	vector

6 entries correct ✓✓✓

4 or 5 entries correct ✓✓

2 or 3 entries correct ✓

(3)

(b)(i)  $mg = EQ$  ✓


$$E \left( = \frac{mg}{Q} = \frac{4.3 \times 10^{-9} \times 9.81}{3.2 \times 10^{-12}} \right) = 1.32 \times 10^4 \text{ (V m}^{-1}\text{)} \quad \checkmark$$

(ii) positive ✓

(3)

(6)

Question 5			
(a)	(i)	$E\left(=\frac{V}{d}\right)=\frac{1400}{15\times 10^{-3}}\checkmark(=9.3\times 10^4\text{ V m}^{-1})$	<b>Q5 Jan 2006</b>
	(ii)	$t\left(=\frac{l}{v}\right)=\frac{30\times 10^{-3}}{3.2\times 10^7}=9.38\times 10^{-10}\text{ s}\checkmark$	
	(iii)	$ma_y = Ee\checkmark$ $a_y = \frac{9.3\times 10^4 \times 1.60\times 10^{-19}}{9.11\times 10^{-31}}\checkmark(=1.64\times 10^{16}\text{ m s}^{-2})$ acceleration is upwards [or towards + plate]✓	
(b)		$v_y(=a_y t)=1.64\times 10^{16}\times 9.38\times 10^{-10}\checkmark(=1.54\times 10^7\text{ m s}^{-1})$ $v=\sqrt{(1.54\times 10^7)^2+(3.2\times 10^7)^2}=3.55\times 10^7\text{ m s}^{-1}\checkmark$ at $\tan^{-1}\left(\frac{1.54}{3.2}\right)=26^\circ$ above the horizontal ✓	<b>5</b>
<b>Total</b>			<b>8</b>

Question 3			
(a)	(i)	force per unit charge ✓ acting on a positive charge ✓	<b>Q3 Jun 2006</b>
	(ii)	vector ✓	
(b)	(i)	$F\left(=\frac{Q_1 Q_2}{4\pi\epsilon_0 r^2}\right)=\frac{4.0\times 10^{-9}\times 8.0\times 10^{-9}}{4\pi\times 8.85\times 10^{-12}\times (80\times 10^{-3})^2}\checkmark$ $=4.5(0)\times 10^{-5}\text{ N}\checkmark$	<b>4</b>
	(ii)	(use of $V=\frac{Q}{4\pi\epsilon_0 r}$ gives) $0=\left(\frac{4.0\times 10^{-9}}{4\pi\epsilon_0 x}\right)-\left(\frac{8.0\times 10^{-9}}{4\pi\epsilon_0(80\times 10^{-3}-x)}\right)$ or $\frac{4}{x}=\frac{8}{80-x}\checkmark$ $x=26.7\text{ mm}\checkmark$	
(c)			<b>3</b>
		correct directions for $E_4$ and $E_8$ ✓ $E_8$ approx twice as long as $E_4$ ✓ correct direction of resultant R shown ✓	
<b>Total</b>			<b>10</b>

<b>Question 4</b>		
(a)	(i)	$E \left( = \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{79 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{12} \times (3.0 \times 10^{-14})^2} \checkmark$ <p style="text-align: center;">(gives <math>E = 1.3 \times 10^{20} \text{ V m}^{-1}</math> (<math>1.26 \times 10^{20}</math>))</p>
	(ii)	$F (= EQ') = 1.26 \times 10^{20} \times 2 \times 1.60 \times 10^{-19} \checkmark$ $= 40 \text{ N (40.3)} \checkmark$
	(iii)	$V \left( = \frac{Q}{4\pi\epsilon_0 r} \right) = \frac{79 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{-12} \times 3.0 \times 10^{-14}} \checkmark$ <p>gives <math>V = 3.8 \times 10^6 \text{ V (or J C}^{-1}\text{)}</math> (<math>3.79 \times 10^6</math>) <math>\checkmark</math></p>
(b)	(i)	kinetic energy $\rightarrow$ electric potential energy $\rightarrow$ kinetic energy $\checkmark$
	(ii)	initial kinetic energy = potential energy at point P $\checkmark$ $= (2e)V \checkmark$ $= 2 \times 1.60 \times 10^{-19} \times 3.79 \times 10^6 = 1.21 \times 10^{-12} \text{ (J)} \checkmark$
		<b>Total</b>
		<b>9</b>

<b>Question 3</b>		
(a)	(i)	force is perpendicular to initial velocity <b>or</b> acts in opposite direction to direction of electric field $\checkmark$ initial velocity component is maintained $\checkmark$
		<b>Q3 Jan 2008</b>
		electron is accelerated in perpendicular direction $\checkmark$ parabolic path $\checkmark$
	(ii)	force is in opposite direction to initial velocity $\checkmark$ electron decelerated $\checkmark$ direction of motion may eventually be reversed $\checkmark$
		<b>max 5</b>

(b)	(i)	$E \left( \frac{V}{d} \right) = \frac{110}{44 \times 10^{-3}} = 2500 \text{ V m}^{-1} \text{ (or NC}^{-1}\text{)} \checkmark$
	(ii)	$F (= EQ) 2500 \times 1.6 \times 10^{-19} = 4.0 \times 10^{-16} \text{ N} \checkmark$
	(iii)	$E_k \text{ gained (= } E_p \text{ lost)} = eV \checkmark$ $= 1.6 \times 10^{-19} \times 110 = 1.7(6) \times 10^{-17} \text{ (J)} \checkmark$ <p><b>[or</b> <math>E_k \text{ gained} = Fd \checkmark</math></p> $= 4.0 \times 10^{-16} \times 44 \times 10^{-3} = 1.7(6) \times 10^{-17} \text{ (J)} \checkmark]$ <p><b>[or use of</b> <math>F = ma</math> and <math>v^2 = u^2 + 2as</math> gives</p> $a = 4.39 \times 10^{14} \text{ (m s}^{-2}\text{)} \text{ and } v^2 = 3.86 \times 10^{13} \text{ (m}^2 \text{ s}^{-2}\text{)} \checkmark$ $E_k \text{ gained} = \frac{1}{2} mv^2 = \frac{1}{2} \times 9.11 \times 10^{-31} \times 3.86 \times 10^{13}$ $= 1.7(6) \times 10^{-17} \text{ J} \checkmark]$
		<b>Total</b>
		<b>9</b>

Question 4				
(a)	(i)	$V = \frac{Q}{4\pi\epsilon_0 r} \text{ and } E_p = eV \checkmark$ $\text{gives } E_p \left( \frac{e^2}{4\pi\epsilon_0 r} \right) = \frac{(1.6 \times 10^{-19})^2}{4\pi \times 8.85 \times 10^{-12} \times 2.0 \times 10^{-15}} \checkmark$ $= 1.15 \times 10^{-13} \text{ J}$	<b>Q4 Jan 2008</b>	<b>5</b>
	(ii)	$2 \times \frac{1}{2} mv^2 = 1.15 \times 10^{-13} \checkmark$ $\text{gives } v^2 = \frac{1.15 \times 10^{-13}}{2 \times 1.67 \times 10^{-27}} \checkmark \therefore v = 5.8(7) \times 10^6 \text{ ms}^{-1} \checkmark$		
(b)	(i)	$\Delta m = 2 \times (2.01355) - (3.01550 + 1.00728) \checkmark (= 4.32 \times 10^{-3} \text{ u})$ $E = 4.32 \times 10^{-3} \times 931.3 = 4.02 \text{ (MeV)} \checkmark$ $= 4.02 \times 10^6 \times 1.6 \times 10^{-19} = 6.4(4) \times 10^{-13} \text{ J} \checkmark$		<b>4</b>
	(ii)	$\text{energy per unit mass} = \frac{6.44 \times 10^{-13}}{4 \times 1.67 \times 10^{-27}}$ $= 9.6(4) \times 10^{13} \text{ (J kg}^{-1}\text{)} \checkmark$ <p style="text-align: center;">[denominator may be <math>2 \times 2.014 \times 1.66 \times 10^{-27}</math>]</p>		
(c)		<p>supply of fuel is almost unlimited (deuterium from sea water) <math>\checkmark</math></p> <p>fewer waste or radioactivity or environmental problems <math>\checkmark</math></p> <p>energy released per unit mass is (generally) greater <math>\checkmark</math></p>		<b>max 2</b>
			<b>Total</b>	<b>11</b>

Question 4				
(a)	(i)	<p>radial straight lines <math>\checkmark</math></p> <p>symmetrical in all directions <math>\checkmark</math></p> <p>directed inwards towards charge <math>\checkmark</math></p> <p>(marks could be taken from diagram)</p>	<b>Q4 Jun 2009</b>	<b>4</b>
	(ii)	<p>line, labelled <b>L</b>, which is a circular arc (or a complete circle) centred on charge <math>\checkmark</math></p>		
(b)	(i)	$E \left( = \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{0.80 \times 10^{-9}}{4\pi \times 8.85 \times 10^{-12} \times (40 \times 10^{-3})^2} \checkmark$ $= 4.50 \times 10^3 \text{ (V m}^{-1}\text{)} \checkmark$		<b>5</b>
	(ii)	<p>point marked at (40, 4.5) <math>\checkmark</math></p> <p>curve of decreasing gradient <math>\checkmark</math></p> <p>correct <math>E \propto (1/r^2)</math> relationship shown by line drawn <math>\checkmark</math></p>		
			<b>Total</b>	<b>9</b>

Question 2		
(a)	graph B ✓	1
(b)	<p style="text-align: right;"><b>Q2 Jan 2010</b></p> <p>for graph A, <math>V \propto \frac{1}{r}</math> ✓</p> <p>for graph B, <math>E \propto \frac{1}{r^2}</math> ✓</p> <p>[if candidate correctly quotes equations for V and E only, with no further explanation in words, allow ✓ only]</p> <p><b>[alternatively</b> allow a fully correct reference to the – and + gradients of graphs A and B respectively in regions R for ✓✓]</p>	2
(c)	<p><math>E = \frac{Q}{4\pi\epsilon_0 r^2}</math> and <math>V = \frac{Q}{4\pi\epsilon_0 r}</math> give <math>E = \frac{V}{r}</math> ✓</p> <p>[no credit for using just <math>E = \frac{V}{d}</math>]</p> <p>∴ potential of the point <math>V = E r = 3.6 \times 10^4 \times 40 \times 10^{-3} = 1.4 \times 10^3 \text{ V}</math> ✓ (1440) (allow <math>\text{J C}^{-1}</math>)</p>	2
	<b>Total</b>	<b>5</b>